IN THE UNITED STATES PATENT AND TRADEMARK OFFICE Atty's Docket No. 101215-72

APPLICANT : Karsten Stein et al.

FILED : Concurrently Herewith

FOR : Method for Multi-fluorescence Detection

PRELIMINARY AMENDMENT

Hon. Assistant Commissioner of Patents Washington, D.C. 20231

Sir:

Prior to examination, please amend the application as follows:

IN THE SPECIFICATION

Page 1, after line 2, please insert --Background of the
Invention--;

Page 3, after line 11, please insert --Summary of the
Invention--;

Page 6, after line 6, please insert --Brief Description of
the Drawings--;

Page 6, after line 15, please insert --Detailed Description

of the Preferred Embodiments -- .

IN THE CLAIMS

Please amend the claims in accordance with the attached marked-up pages. A clean copy of the amended claims is also enclosed.

REMARKS

The above amendments were made to place the application into proper United States Patent Format.

Respectfully Submitted,

Bruce 3. Londa

Attorney for Applicant

Norris, McLaughlin & Marcus P.A.

220 East 42nd Street, 30th Floor

New York, N.Y. 10017

Telephone: (212)808-0700

Telecopier: (212)808-0844

- 1. Method for the multi-fluorescence detection of fluorophores by means of a simultaneous measurement of the decay time of the fluorescences, where the excitation wave lengths for the individual fluorophores, delayed through an optical delay (4) in the range of sub-nanoseconds to some milliseconds, are conducted to the objects of examination (7) so that the fluorescences can be excitated and detected one after the other and where, for he differentiation between at least two fluorophores in addition to their spectral characteristics, the decay behaviour of the fluorescence processes is examined by the displacement of electronic gates in the nanosecond range along a timing axis.
- 2. Method for the multi-fluorescence detection of
 fluorephores by means of a simultaneous measurement of the decay
 time of the fluorescences, where the excitation wave lengths for
 the individual fluorephores, delayed through an optical delay
 (4) in the range of sub nanoseconds to some milliseconds, are
 conducted to the objects of examination (7) so that the
 fluorescences can be excitated and detected one after the other.
- 3. <u>(amended) Method A method</u> for the multi-fluorescence detection of fluorophores by means of a simultaneous measurement

of the decay time of the fluorescences where, for the differentiation between at least two fluorophores in addition to their spectral characteristics, the decay behaviour of the fluorescence processes is examined by the displacement of electronic gates in the nanosecond range along a timing axis.

- 4. <u>(amended) Method The method according to Claim 1 and 3,</u> wherein the delay (4) is formed by light wave conductors.
- 5. <u>(amended) Method The method</u> according to Claim 2 and 3, wherein the electronic time gate is positioned in the maximum of the timing pattern of the life duration of the fluorescence signal, in order to selectively detect fast decaying fluorescence processes.
- 6. <u>(amended)</u> Method The method according to Claim 2 and 3, wherein the electronic time gate is positioned in the fade-out of the timing pattern of the life duration of the fluorescence signal, in order to selectively detect slow decaying fluorescence processes.

- 7. <u>(amended)</u> Method The method according to the Claims 1 to Claim 3, wherein several different fluorescence colouring materials are detected in the liquid chromatography.
- 8. <u>(amended)</u> Method The method according to the Claims 1 to Claim 3, wherein fluorescence colouring materials are detected in multi-well plates.
- 9. <u>(amended) Method The method</u> according to the Claims 1 to Claim 3, wherein a multiple fluorescence detection is carried out on living/dead tissue.
- 10. <u>(amended)</u> Method The method according to the Claims 1 to Claim 3, wherein a multi fluorescence detection is carried out on planar, particular, fibrillar carriers such as DNA-/protein-chip.
- 11. <u>(amended)</u> <u>Method</u> <u>The method</u> according to <u>the Claims 1</u> to <u>Claim 3</u>, wherein the method is image-rendering and the detector is a camera.

- 12. <u>(amended)</u> <u>Method The method according to the Claims 1</u>
 to Claim 3, wherein a multiple fluorescence detection and an end-point determination is carried out during the PCR, particularly quantitative and multiplex PCR.
- 13. <u>(amended)</u> <u>Method_The method_according to the Claims 1</u>
 to <u>Claim 3</u>, wherein several fluorescence colouring materials are detected in electrophoresis gels, electrophoresis capillaries and electrophoresis blots.
- 14. (new) A method for the multi-fluorescence detection of fluorophores by means of a simultaneous measurement of the decay time of the fluorescences, where the excitation wave lengths for the individual fluorophores, delayed through an optical delay (4) in the range of sub-nanoseconds to some milliseconds, are conducted to the objects of examination (7) so that the fluorescences can be excitated and detected one after the other.
- 15. (new) The method according to claim 14, wherein for the differentiation between at least two fluorophores in addition to their spectral characteristics, the decay behaviour of the

fluorescence processes is examined by the displacement of electronic gates in the nanosecond range along a timing axis.

- 16. (new) The method according to Claim 14, wherein the delay (4) is formed by light wave conductors.
- 17. (new) The method according to Claim 14, wherein the electronic time gate is positioned in the maximum of the timing pattern of the life duration of the fluorescence signal, in order to selectively detect fast decaying fluorescence processes.
- 18. (new) The method according to Claim 14, wherein the electronic time gate is positioned in the fade-out of the timing pattern of the life duration of the fluorescence signal, in order to selectively detect slow decaying fluorescence processes.
- 19. (new) The method according to the Claim 14, wherein several different fluorescence colouring materials are detected in the liquid chromatography.

- 20. (new) The method according to the Claim 14, wherein fluorescence colouring materials are detected in multi-well plates.
- 21. (new) The method according to the Claim 14, wherein a multiple fluorescence detection is carried out on living/dead tissue.
- 22. (new) The method according to the Claim 14, wherein a multi fluorescence detection is carried out on planar, particular, fibrillar carriers such as DNA-/protein-chip.
- 23. (new) The method according to the Claim 14, wherein the method is image-rendering and the detector is a camera.
- 24. (new) The method according to the Claim 14, wherein a multiple fluorescence detection and an end-point determination is carried out during the PCR, particularly quantitative and multiplex PCR.
- 25. (new) The method according to the Claim 14, wherein several fluorescence colouring materials are detected in

electrophoresis gels, electrophoresis capillaries and electrophoresis blots.

- 3. (amended) A method for the multi-fluorescence detection of fluorophores by means of a simultaneous measurement of the decay time of the fluorescences where, for the differentiation between at least two fluorophores in addition to their spectral characteristics, the decay behaviour of the fluorescence processes is examined by the displacement of electronic gates in the nanosecond range along a timing axis.
- 4. (amended) The method according to Claim 3, wherein the delay (4) is formed by light wave conductors.
- 5. (amended) The method according to Claim 3, wherein the electronic time gate is positioned in the maximum of the timing pattern of the life duration of the fluorescence signal, in order to selectively detect fast decaying fluorescence processes.
- 6. (amended) The method according to Claim 3, wherein the electronic time gate is positioned in the fade-out of the timing pattern of the life duration of the fluorescence signal, in

order to selectively detect slow decaying fluorescence processes.

- 7. (amended) The method according to Claim 3, wherein several different fluorescence colouring materials are detected in the liquid chromatography.
- 8. (amended) The method according to Claim 3, wherein fluorescence colouring materials are detected in multi-well plates.
- 9. (amended) The method according to Claim 3, wherein a multiple fluorescence detection is carried out on living/dead tissue.
- 10. (amended) The method according to Claim 3, wherein a multi fluorescence detection is carried out on planar, particular, fibrillar carriers such as DNA-/protein-chip.
- 11. (amended) The method according to Claim 3, wherein the method is image-rendering and the detector is a camera.

- 12. (amended) The method according to Claim 3, wherein a multiple fluorescence detection and an end-point determination is carried out during the PCR, particularly quantitative and multiplex PCR.
- 13. (amended) The method according to Claim 3, wherein several fluorescence colouring materials are detected in electrophoresis gels, electrophoresis capillaries and electrophoresis blots.
- 14. (new) A method for the multi-fluorescence detection of fluorophores by means of a simultaneous measurement of the decay time of the fluorescences, where the excitation wave lengths for the individual fluorophores, delayed through an optical delay (4) in the range of sub-nanoseconds to some milliseconds, are conducted to the objects of examination (7) so that the fluorescences can be excitated and detected one after the other.
- 15. (new) The method according to claim 14, wherein for the differentiation between at least two fluorophores in addition to their spectral characteristics, the decay behaviour of the

fluorescence processes is examined by the displacement of electronic gates in the nanosecond range along a timing axis.

- 16. (new) The method according to Claim 14, wherein the delay (4) is formed by light wave conductors.
- 17. (new) The method according to Claim 14, wherein the electronic time gate is positioned in the maximum of the timing pattern of the life duration of the fluorescence signal, in order to selectively detect fast decaying fluorescence processes.
- 18. (new) The method according to Claim 14, wherein the electronic time gate is positioned in the fade-out of the timing pattern of the life duration of the fluorescence signal, in order to selectively detect slow decaying fluorescence processes.
- 19. (new) The method according to the Claim 14, wherein several different fluorescence colouring materials are detected in the liquid chromatography.

- 20. (new) The method according to the Claim 14, wherein fluorescence colouring materials are detected in multi-well plates.
- 21. (new) The method according to the Claim 14, wherein a multiple fluorescence detection is carried out on living/dead tissue.
- 22. (new) The method according to the Claim 14, wherein a multi fluorescence detection is carried out on planar, particular, fibrillar carriers such as DNA-/protein-chip.
- 23. (new) The method according to the Claim 14, wherein the method is image-rendering and the detector is a camera.
- 24. (new) The method according to the Claim 14, wherein a multiple fluorescence detection and an end-point determination is carried out during the PCR, particularly quantitative and multiplex PCR.
- 25. (new) The method according to the Claim 14, wherein several fluorescence colouring materials are detected in

electrophoresis gels, electrophoresis capillaries and electrophoresis blots.